

ASTHMA ACTION PLAN

Take this ASTHMA ACTION PLAN with you when you visit your doctor



NAME	DOCTOR'S CONTACT DETAILS	EMERGENCY CONTACT DETAILS
DATE		Name
NEXT ASTHMA CHECK-UP DUE		Phone
.....		Relationship

WHEN WELL *Asthma under control (almost no symptoms)* **ALWAYS CARRY YOUR RELIEVER WITH YOU**

Peak flow* (if used) above:

Your preventer is: (NAME & STRENGTH)

Take puffs/tablets times every day

Use a spacer with your inhaler

Your reliever is: (NAME)

Take puffs

When: You have symptoms like wheezing, coughing or shortness of breath

Use a spacer with your inhaler

OTHER INSTRUCTIONS
(e.g. other medicines, trigger avoidance, what to do before exercise)

.....

.....

.....

.....

WHEN NOT WELL *Asthma getting worse (needing more reliever than usual, having more symptoms than usual, waking up with asthma, asthma is interfering with usual activities)*

Peak flow* (if used) between and

Keep taking preventer: (NAME & STRENGTH)

Take puffs/tablets times every day

.....

Use a spacer with your inhaler

Your reliever is: (NAME)

Take puffs

.....

Use a spacer with your inhaler

OTHER INSTRUCTIONS Contact your doctor
(e.g. other medicines, when to stop taking extra medicines)

.....

.....

.....

.....

IF SYMPTOMS GET WORSE *Severe asthma flare-up/attack (needing reliever again within 3 hours, increasing difficulty breathing, waking often at night with asthma symptoms)*

Peak flow* (if used) between and

Keep taking preventer: (NAME & STRENGTH)

Take puffs/tablets times every day

.....

Use a spacer with your inhaler

Your reliever is: (NAME)

Take puffs

.....

Use a spacer with your inhaler

OTHER INSTRUCTIONS Contact your doctor today
(e.g. other medicines, when to stop taking extra medicines)

Prednisolone/prednisone:

Take each morning for days

.....

.....

.....

DANGER SIGNS *Asthma emergency (severe breathing problems, symptoms get worse very quickly, reliever has little or no effect)*

Peak flow (if used) below:

DIAL 000 FOR AMBULANCE

Call an ambulance immediately
Say that this is an asthma emergency
Keep taking reliever as often as needed

Use your adrenaline autoinjector (EpiPen or Anapen)

National Asthma Council Australia
leading the attack against asthma

nationalasthma.org.au

* Peak flow not recommended for children under 12 years.

ASTHMA ACTION PLAN

what to look out for



WHEN WELL



THIS MEANS:

- you have no night-time wheezing, coughing or chest tightness
- you only occasionally have wheezing, coughing or chest tightness during the day
- you need reliever medication only occasionally or before exercise
- you can do your usual activities without getting asthma symptoms

WHEN NOT WELL



THIS MEANS ANY ONE OF THESE:

- you have night-time wheezing, coughing or chest tightness
- you have morning asthma symptoms when you wake up
- you need to take your reliever more than usual
- your asthma is interfering with your usual activities

THIS IS AN ASTHMA FLARE-UP

IF SYMPTOMS GET WORSE



THIS MEANS:

- you have increasing wheezing, cough, chest tightness or shortness of breath
- you are waking often at night with asthma symptoms
- you need to use your reliever again within 3 hours

THIS IS A SEVERE ASTHMA ATTACK (SEVERE FLARE-UP)

DANGER SIGNS



THIS MEANS:

- your symptoms get worse very quickly
- you have severe shortness of breath, can't speak comfortably or lips look blue
- you get little or no relief from your reliever inhaler

**CALL AN AMBULANCE IMMEDIATELY: DIAL 000
SAY THIS IS AN ASTHMA EMERGENCY**

**DIAL 000 FOR
AMBULANCE**

ASTHMA MEDICINES

PREVENTERS

Your preventer medicine reduces inflammation, swelling and mucus in the airways of your lungs. Preventers need to be taken **every day**, even when you are well.

Some preventer inhalers contain 2 medicines to help control your asthma (combination inhalers).

RELIEVERS

Your reliever medicine works quickly to make breathing easier by making the airways wider.

Always carry your reliever with you – it is essential for first aid. Do not use your preventer inhaler for quick relief of asthma symptoms unless your doctor has told you to do this.

To order more Asthma Action Plans visit the National Asthma Council website. A range of action plans are available on the website – please use the one that best suits your patient.

nationalasthma.org.au

Developed by the National Asthma Council Australia and supported by GSK Australia.

National Asthma Council Australia retained editorial control. © 2015

National Asthma
Council Australia
leading the attack against asthma

nationalasthma.org.au